

Top Quark Physics at the Tevatron

Shinhong Kim

University of Tsukuba

International Workshop on Linear Colliders LCWS99, April 30, 1999

- Tevatron and CDF Detector in Run2
- Top Quark Physics, present
- Top Quark Physics, future



Tevatron and CDF Detector in RUN2

RUN2(will start in April 2000)

- Accelerator Upgrade
 - CM Energy: 1.8TeV \rightarrow 2.0TeV
 - Integrated Luminosity: 0.1 fb⁻¹ \rightarrow 2 fb⁻¹
 - Instantaneous Luminosity: 2.5 \times 10³¹ \rightarrow 5 \times 10³¹ \sim 2 \times 10³²
 - Number of Bunches: 6 \rightarrow 36 \sim 108
 - Bunch Period: 3.5 μ s \rightarrow 396 \sim 132 ns
- Detector Upgrade (CDF II detector, Technical Design Report Nov. 1996)
 - Quick response to the shorter bunch period
 - * CTC \rightarrow COT
 - * Gas Calorimeter \rightarrow Scintillator Calorimeter
 - Large coverage of SVX II (twice larger than SVX')

Collider Detector at Fermilab (CDF)

CDF Detector Evolution



History of the Top Quark Search

- 1977-1994: **A fine collection of null results**

- April, 1994: **First Evidence**

- **Phys. Rev. D50, 2966 (1994) CDF**

- 15 events on a background of 6.0

- 2.8 σ excess

- $M_{top} = 174 \pm 17 \text{ GeV}/c^2$

- $\sigma_{t\bar{t}} = 13.9_{-4.8}^{+6.1} \text{ pb}$

- February, 1995: **Confirmation**

- **PRL 74, 2626 (1995) CDF**

- 4.8 σ excess

- $M_{top} = 176 \pm 8(\text{stat}) \pm 10(\text{syst}) \text{ GeV}/c^2$

- $\sigma_{t\bar{t}} = 6.8_{-2.4}^{+3.6} \text{ pb}$

- **PRL 74, 2632 (1995) D0**


- 4.6 σ excess

- $M_{top} = 199_{-21}^{+19} (\text{stat}) + 14_{-21}(\text{syst}) \text{ GeV}/c^2$

- $\sigma_{t\bar{t}} = 6.4 \pm 2.2 \text{ pb}$

Top Production at the Tevatron

Top quarks are predominantly produced in pairs by the process $p\bar{p} \rightarrow t\bar{t}$.




In Tevatron Run I,

- Integrated Luminosity exceeded 100 pb^{-1}
- over 5×10^{12} $p\bar{p}$ collisions
- **≈ 500 $t\bar{t}$ pairs produced.**

$$\frac{\sigma_{t\bar{t}}}{\sigma_{\text{inel}}} \sim 10^{-9}, \quad \frac{\sigma_{t\bar{t}}}{\sigma_{\text{W}}} \sim 10^{-3}$$

Top Quark Decay Signatures



We categorize top decays by how the two W bosons decay.

- Both W 's Decay $W \rightarrow l\nu$ (Dilepton Channel)

Final State: $l^+\nu l^-\nu b\bar{b}$ ($l:e$ or μ ; 5%)


- One W Decay $W \rightarrow l\nu$ (Lepton+Jets Channel)

Final State: $l^+\nu q\bar{q}' b\bar{b}$ ($l:e$ or μ ; 30%)

- Both W 's Decay $W \rightarrow q\bar{q}'$ (All Hadronic Channel)

Final State: $q\bar{q}' q\bar{q}' b\bar{b}$ (44%)

Lepton + Jets Channels



Signature:

- One isolated high P_T lepton (e or μ)
- Missing Energy (E_T)
- 4 or more jets, 2 of which are from b-quarks

Dominant Backgrounds:

- $p\bar{p} \rightarrow W + jets$
- QCD background (Fake leptons)

Two Distinct Differences between $t\bar{t}$ and $W+jets$


- $t\bar{t}$ events always contain b quarks, $W+jet$ events usually do not.
- The jets in $W+jet$ events tend to be softer than in $t\bar{t}$ (low E_t)

Need to further suppress backgrounds either with:


- Topological/kinematic requirements:
 - Aplanarity
 - $H_T = E_T(\text{leptons}) + \cancel{E}_T + \sum E_t(\text{jets})$
 - Form likelihood ($t\bar{t}$ vs. Bkgd) based on Jet E_t 's
- Tag b -quarks using semileptonic decays (SLT b -tag)

$$b \rightarrow eX, \quad b \rightarrow \mu X \quad (20\%)$$

- Tag b -quarks using Displaced vertex (SVX b -tag)



CDF W+Jet Events



The final numbers are in

We have (CDF and D0 Combined):

- ≈ 13 Dileptons
- ≈ 60 Lepton+Jets
- ≈ 60 All Hadronic

What do we do next ?


- Production Properties ($\sigma_{t\bar{t}}$)
- Decay Properties (V_{tb})
- Top Quark Mass
- Search for new physics (Rare decays, $M_{t\bar{t}}$)

$\sigma_{t\bar{t}}$ Measurements

$$\sigma = \frac{N_{obs} - N_{bkg}}{A\mathcal{L}}$$


Goal: Determine the top cross section

- as accurately as possible
- in as many different decay channels as possible as a check of top decay and compare it to theoretical prediction.



Top Quark Mass in Lepton + Jet Channel at CDF

76 Events




$$M_{\text{top}} = 175.9 \pm 4.8(\text{stat}) \pm 5.3(\text{syst}) \text{ GeV}/c^2$$

Combining this with the measurements in the dilepton and all-hadronic channels,

$$M_{\text{top}} = 176.0 \pm 6.5 \text{ GeV}/c^2 \text{ (CDF)}$$


Comparison between CDF and D0



CDF + D0 Combined Top Mass

$$\begin{aligned} M_{top} &= 174.3 \pm 3.2(\text{stat}) \pm 4.0(\text{syst}) \text{ GeV}/c^2 \\ &= 174.3 \pm 5.1 \text{ GeV}/c^2 \end{aligned}$$

M_W vs M_T



Higgs Mass Constraint

From M_{top} (CDF,D0), M_W (CDF,D0,LEP2) and other electroweak results at LEP and SLC


- $M_H = 76^{+85}_{-47} \text{ GeV}/c^2$
- $M_H < 262 \text{ GeV}/c^2$ at 95% C. L.

[ref] LEP Electroweak Working Group, CERN EP/99-15.

Search for single top quark production


s-channel W^* process

$\sigma_{\text{theory}} = 0.73 \text{ pb}$
Signal is $W+b+\bar{b}$



W-gluon fusion process

$\sigma_{\text{theory}} = 1.70 \text{ pb}$
Signal is $W+b+q$



- Event selection

- Isolated lepton $E_T \geq 20 \text{ GeV}$
- Missing $E_T \geq 20 \text{ GeV}$
- Exactly two jets with uncorrected $E_T \geq 15 \text{ GeV}$ and $|\eta| \leq 2$.
- Standard Z and top dilepton removal

W^* signal search:

at least one *b*-tagged jet

W-gluon fusion signal search:



only one *b*-tagged jet + top mass window cut

W-gluon fusion single top search

Candidate : 15 events
 Expected background ($Wb\bar{b}, t\bar{t} \dots$) : 12.9 ± 2.1 events
 Expected signal : 1.2 ± 0.3 events

Perform a likelihood fit of $Q \times \eta$ distribution:

Fitted background : 13.1 ± 1.9 events
 Fitted signal : $1.4^{+4.2}_{-3.4}$ events



Q : lepton charge (± 1)



η : untagged jet pseudo-rapidity

W^* single top search

Candidate : 42 events
Expected background : 31.3 ± 4.7
Expected signal : 1.0 ± 0.3

Perform a likelihood fit of reconstructed top mass distribution:

Fitted background : $33.0^{+4.4}_{-4.3}$
Fitted signal : $6.6^{+7.3}_{-6.5}$



Top Quark Physics in RUN2

- Accelerator Upgrade
 - CM Energy:
 - * 1.8TeV \rightarrow 2.0TeV ($\sigma_{t\bar{t}}$: \times 1.4)
 - Integrated Luminosity:
 - * 0.1 fb $^{-1}$ \rightarrow 2 fb $^{-1}$ (\times 20)
- Detector Upgrade (CDF)
 - Acceptance for $t\bar{t} \rightarrow W(\rightarrow \ell\nu)_+ \geq 3$ jets:
 - * 8.7% \rightarrow 10% (\times 1.15)
 - b -tagging efficiency for $t\bar{t} \rightarrow W(\rightarrow \ell\nu)_+ \geq 3$ jets:
 - * 52% \rightarrow 86% (\times 1.65)


From the above, we have 50 times higher rate of $t\bar{t} \rightarrow W(\rightarrow \ell\nu)_+ \geq 3$ jet events with one b -tagged jet in RUN2 (2fb $^{-1}$).

Number of $t\bar{t}$ events in RUN2(CDF)

- $W(\rightarrow \ell\nu)_+ \geq 4$ jets (1 b -tag): \sim 1,000
- $W(\rightarrow \ell\nu)_+ \geq 4$ jets (2 b -tags): \sim 500

CDF RUN2 (2fb⁻¹)

- Top Quark Mass: $\Delta M_{top} \sim 3\text{GeV}/c^2$
- $\sigma_{t\bar{t}}$: $\Delta\sigma/\sigma \sim 9\%$
- σ_{top} from $q\bar{q} \rightarrow W^* \rightarrow t\bar{b}$ and $gW^* \rightarrow t\bar{b}$: $\Delta, /, \sim 25\%$
- V_{tb} : $\Delta|V_{tb}|/|V_{tb}| \sim 13\%$
- New Particle Search such as $t \rightarrow H^+b$ or $X \rightarrow t\bar{t}$.



From M_{top} (Tevatron RUN2), M_W (Tevatron RUN2,LEP II) and other electroweak results at LEP and SLC,

$$\Delta M_H/M_H \sim 30\%$$

[Ref] TEV2000 Rroup Report FERMILAB-PUB-96/082 and Light Higgs Working Group Report at SNOWMASS 96.